Copyright

Submission of a manuscript implies: that the work described has not been published before (except in the form of an abstract); that it is not under consideration for publication elsewhere; that its publication has been approved by all coauthors, if any, as well as by the responsible authorities at the institute where the work has been carried out; that, if and when the manuscript is accepted for publication, the authors agree to automatic transfer of the copyright to the publisher and that the manuscript will not be published elsewhere in any language without the consent of the copyright holders.

All articles published in this journal are protected by copyright, which covers the exclusive rights to reproduce and distribute the article (e. g., as offprints), all translation rights as well as the rights to publish the article in any electronic form. No material published in this journal may be reproduced photographically or stored on microfilm, in electronic data bases, video disks, etc., without first obtaining written permission from the publisher.

The use of general descriptive names, trade names, trademarks, etc., in this publication, even if not specifically identified, does not imply that these names are not protected by the relevant laws and regulations.

While the advice and information in this journal is believed to be true and accurate at the date of its going to press, neither the authors, the editors, nor the publisher can accept any legal responsibility for any errors or omissions that may be made. The publisher makes no warranty, express or implied, with respect to the material contained herein.

Special regulations for photocopies in the USA. Photocopies may be made for personal or in-house use beyond the limitations stipulated under Section 107 or 108 of U.S. Copyright Law, provided a fee is paid. All fees should be paid to the Copyright Clearance Center, Inc., 21 Congress Street, Salem, MA 01970, USA, stating the ISSN 0012-186X, the volume, and the first and last page numbers of each article copied. The copyright owner's consent does not include copying for general distribution, promotion, new works, or resale. In these cases, specific written permission must first be obtained from the publisher.

Springer-Verlag Berlin Heidelberg New York London Paris Tokyo Hong Kong Barcelona Budapest Printed in Germany by aprinta Wemding
© Springer-Verlag Berlin Heidelberg 2001

Contents

Editorial

Tomkin G. H.: Diabetic vascular disease and the rising star of Protein Kinase C 657

Waldhäusl W. K.: Finally we have arrived in a new millennium 1

Reviews

- Bonora E., Muggeo M.: Postprandial blood glucose as a risk factor for cardiovascular disease in Type II diabetes: the epidemiological evidence 2107
- Cameron N. E., Eaton S. E. M., Cotter M. A., Tesfaye S.: Vascular factors and metabolic interactions in the pathogenesis of diabetic neuropathy 1973
- Chatenoud L.: Restoration of self-tolerance is a feasible approach to control ongoing beta-cell specific autoreactivity: its relevance for treatment in established diabetes and islet transplantation 521
- Cooper M. E.: Interaction of metabolic and haemodynamic factors in mediating experimental diabetic nephropathy 1957

- Dalgaard L.T., Pedersen O: Uncoupling proteins: functional characteristics and role in the pathogenesis of obesity and Type II diabetes 946
- Di Mario U., Pugliese G: 15th Golgi lecture: from hyperglycaemia to the dysregulation of vascular remodelling in diabetes 674
- Edlund H.: Factors controlling pancreatic cell differentiation and function
- Eizirik D. L., Mandrup-Poulsen T.: A choice of death – the signal-transduction of immune-mediated beta-cell apoptosis 2115
- Gale É. A. M., Gillespie K. M.: Diabetes and gender 3
- Hu F. B., van Dam R. M., Liu S: Diet and risk of Type II diabetes: the role of types of fat and carbohydrate 805
- Idris I., Gray S., Donnelly R.: Protein kinase C activation: isozyme-specific effects on metabolism and cardiovascular complications in diabetes 659
- Jun H. S., Yoon J. W.: The role of viruses in Type I diabetes: two distinct cellular and molecular pathogenic mechanisms of virus-induced diabetes in animals 271
- Lorenzi M., Gerhardinger C.: Early cellular and molecular changes induced by diabetes in the retina 791

- Malaisse W.J.: On the track to the betacell 393
- McKinnon C. M., Docherty K.: Pancreatic duodenal homeobox-1, PDX-1, a major regulator of beta cell identity and function 1203
- Pratley R. E., Weyer C.: The role of impaired early insulin secretion in the pathogenesis of Type II diabetes mellitus 929
- Singh R., Barden A., Mori T., Beilin L.: Advanced glycation end-products: a review 129
- Soria B., Skoudy A., Martín F.: From stem cells to beta cells: new strategies in cell therapy of diabetes mellitus 407
- Stacher G: Diabetes mellitus and the stomach 1080

Articles

- Ablamunits V., Bridgett M., Duffy T., Haag F., Nissen M., Koch-Nolte F., Leiter H.: Changing patterns of cell surface mono (ADP-ribosyl) transferase antigen ART2.2 on resting versus cytopathically-activated T cells in NOD/Lt mice 848
- ADVANCE Management Committee: Study Rationale and Design of AD-

Copyright

Submission of a manuscript implies: that the work described has not been published before (except in the form of an abstract); that it is not under consideration for publication elsewhere; that its publication has been approved by all coauthors, if any, as well as by the responsible authorities at the institute where the work has been carried out; that, if and when the manuscript is accepted for publication, the authors agree to automatic transfer of the copyright to the publisher and that the manuscript will not be published elsewhere in any language without the consent of the copyright holders.

All articles published in this journal are protected by copyright, which covers the exclusive rights to reproduce and distribute the article (e. g., as offprints), all translation rights as well as the rights to publish the article in any electronic form. No material published in this journal may be reproduced photographically or stored on microfilm, in electronic data bases, video disks, etc., without first obtaining written permission from the publisher.

The use of general descriptive names, trade names, trademarks, etc., in this publication, even if not specifically identified, does not imply that these names are not protected by the relevant laws and regulations.

While the advice and information in this journal is believed to be true and accurate at the date of its going to press, neither the authors, the editors, nor the publisher can accept any legal responsibility for any errors or omissions that may be made. The publisher makes no warranty, express or implied, with respect to the material contained herein.

Special regulations for photocopies in the USA. Photocopies may be made for personal or in-house use beyond the limitations stipulated under Section 107 or 108 of U.S. Copyright Law, provided a fee is paid. All fees should be paid to the Copyright Clearance Center, Inc., 21 Congress Street, Salem, MA 01970, USA, stating the ISSN 0012-186X, the volume, and the first and last page numbers of each article copied. The copyright owner's consent does not include copying for general distribution, promotion, new works, or resale. In these cases, specific written permission must first be obtained from the publisher.

Springer-Verlag Berlin Heidelberg New York London Paris Tokyo Hong Kong Barcelona Budapest Printed in Germany by aprinta Wemding
© Springer-Verlag Berlin Heidelberg 2001

Contents

Editorial

Tomkin G. H.: Diabetic vascular disease and the rising star of Protein Kinase C 657

Waldhäusl W. K.: Finally we have arrived in a new millennium 1

Reviews

- Bonora E., Muggeo M.: Postprandial blood glucose as a risk factor for cardiovascular disease in Type II diabetes: the epidemiological evidence 2107
- Cameron N. E., Eaton S. E. M., Cotter M. A., Tesfaye S.: Vascular factors and metabolic interactions in the pathogenesis of diabetic neuropathy 1973
- Chatenoud L.: Restoration of self-tolerance is a feasible approach to control ongoing beta-cell specific autoreactivity: its relevance for treatment in established diabetes and islet transplantation 521
- Cooper M. E.: Interaction of metabolic and haemodynamic factors in mediating experimental diabetic nephropathy 1957

- Dalgaard L.T., Pedersen O: Uncoupling proteins: functional characteristics and role in the pathogenesis of obesity and Type II diabetes 946
- Di Mario U., Pugliese G: 15th Golgi lecture: from hyperglycaemia to the dysregulation of vascular remodelling in diabetes 674
- Edlund H.: Factors controlling pancreatic cell differentiation and function
- Eizirik D. L., Mandrup-Poulsen T.: A choice of death – the signal-transduction of immune-mediated beta-cell apoptosis 2115
- Gale É. A. M., Gillespie K. M.: Diabetes and gender 3
- Hu F. B., van Dam R. M., Liu S: Diet and risk of Type II diabetes: the role of types of fat and carbohydrate 805
- Idris I., Gray S., Donnelly R.: Protein kinase C activation: isozyme-specific effects on metabolism and cardiovascular complications in diabetes 659
- Jun H. S., Yoon J. W.: The role of viruses in Type I diabetes: two distinct cellular and molecular pathogenic mechanisms of virus-induced diabetes in animals 271
- Lorenzi M., Gerhardinger C.: Early cellular and molecular changes induced by diabetes in the retina 791

- Malaisse W.J.: On the track to the betacell 393
- McKinnon C. M., Docherty K.: Pancreatic duodenal homeobox-1, PDX-1, a major regulator of beta cell identity and function 1203
- Pratley R. E., Weyer C.: The role of impaired early insulin secretion in the pathogenesis of Type II diabetes mellitus 929
- Singh R., Barden A., Mori T., Beilin L.: Advanced glycation end-products: a review 129
- Soria B., Skoudy A., Martín F.: From stem cells to beta cells: new strategies in cell therapy of diabetes mellitus 407
- Stacher G: Diabetes mellitus and the stomach 1080

Articles

- Ablamunits V., Bridgett M., Duffy T., Haag F., Nissen M., Koch-Nolte F., Leiter H.: Changing patterns of cell surface mono (ADP-ribosyl) transferase antigen ART2.2 on resting versus cytopathically-activated T cells in NOD/Lt mice 848
- ADVANCE Management Committee: Study Rationale and Design of AD-

VANCE: Action in Diabetes and Vascular disease – preterax and diamicron MR controlled evaluation (Short com-

munication) 1118

Ahn J. D., Morishita R., Kaneda Y., Lee K. U., Park J. Y., Jeon Y. J., Song H. S., Lee I. K.: Transcription factor decoy for activator protein-1 (AP-1) inhibits high glucose- and angiotensin II-induced type 1 plasminogen activator inhibitor (PAI-1) gene expression in cultured human vascular smooth muscle cells 713

Ahrén, B., Larsson H.: Impaired glucose tolerance (IGT) is associated with reduced insulin-induced suppression of glucagon concentrations 1998

Aoki M., Morishita R., Hayashi S., Jo N., Matsumoto K., Nakamura T., Kaneda Y., Ogihara T.: Inhibition of neointimal formation after balloon injury by cilostazol, accompanied by improvement of endothelial dysfunction and induction of hepatocyte growth factor in rat diabetes model 1034

Assert R., Scherk G., Bumbure A., Pirags V., Schatz H., Pfeiffer A. F. H.:
Regulation of protein kinase C by short term hyperglycaemia in human platelets in vivo and in vitro 188

Bagust A., Hopkinson P.K., Maier W., Currie C.J.: An economic model of the long-term health care burden of Type II diabetes 2140

Baird J., Osmond C., MacGregor A., Snieder H., Hales C. N., Phillips D. I. W.: Testing the fetal origins hypothesis in twins: the Birmingham twin study 33

Barberà A., Gomis R.R., Prats N., Rodríguez-Gil J. E., Domingo M., Gomis R., Guinovart J.J.: Tungstate is an effective antidiabetic agent in streptozotocin-induced diabetic rats: a longterm study 507

Barbieri M., Ragno E., Benvenuti E., Zito G.A., Corsi A., Ferrucci L., Paolisso G.: New aspects of the insulin resistance syndrome: impact on haematological parameters 1232

Bardoux P., Ahloulay M., Le Maout S., Bankir L., Trinh-Trang-Tan M.-M.: Aquaporin-2 and urea transporter-A1 are up-regulated in rats with Type I diabetes mellitus 637

Baroni M. G., Arca M., Sentinelli F., Buzzetti R., Capici F., Lovari S., Vitale M., Romeo S., Di Mario U.: The G972R variant of the Insulin Receptor Substrate-1 (IRS-1) gene, body fat distribution and insulin-resistance 367

Benhamou P. Y., Oberholzer J., Toso C., Kessler L., Penfornis A., Bayle F., Thivolet C., Martin X., Ris F., Badet L., Colin C., Morel P on behalf of the GRAGIL group: Human islet transplantation network for the treatment of Type I diabetes: first data from the Swiss-French GRAGIL consortium (1999–2000) 859

Bergman M.-L., Penha-Gonçalves C., Lejon K., Holmberg D.: Low rate of proliferation in immature thymocytes of the non-obese diabetic mouse maps to the *Idd6* diabetes susceptibility region 1054

Bertelli E., Regoli M., Orazioli D., Bendayan M.: Association between islets of Langerhans and pancreatic ductal system in adult rat. Where endocrine and exocrine meet together? 575

Bertelsen M., Änggård E.E., Carrier M.J.: Oxidative stress impairs insulin internalization in endothelial cells in vitro 605

Bertuglia S., Giusti A., Fedele S., Picano E.: Glucose-insulin-potassium treatment in combination with dipyridamole inhibits ischaemia-reperfusion-induced damage 2165

Beyer U., Schäfer D., Thomas A., Aulich H., Haueter U., Reihl B., Ehwald R.: Recording of subcutaneous glucose dynamics by a viscometric affinity sensor (Rapid communication) 416

Biessels G.J., Braun K.P.J., de Graaf R.A., van Eijsden P., Gispen W.H., Nicolay K.: Cerebral metabolism in streptozotocin-diabetic rats: an in vivo magnetic resonance spectroscopy study 346

Blaak E. E., Glatz J. F. C., Saris W. H. M.: Increase in skeletal muscle fatty acid binding protein (FABPC) content is directly related to weight loss and to changes in fat oxidation following a very low calorie diet (Short communication) 2013

Bo S., Menato G., Lezo A., Signorile A., Bardelli C., Michieli FDe, Massobrio M., Pagano G.: Dietary fat and gestational hyperglycaemia 972

Bonnet F., Cooper M. E., Kawachi H., Allen T. J., Boner G., Cao Z.: Irbesartan normalises the deficiency in glomerular nephrin expression in a model of diabetes and hypertension 874

Borrebaek J., Prydz K., Fjeldstad K., Vuong T.T., Berg T.J., Holkov C., Kolset S. O.: The AGE product N^c-(carboxymethyl)lysine serum albumin is a modulator of proteoglycan expression in polarized cultured kidney epithelial cells 488

Boulton A.J. M., Selam J.-L., Sweeney M., Ziegler D.: Sildenafil citrate for the treatment of erectile dysfunction in men with Type II diabetes mellitus 1296

Boutin P., Vasseur F., Samson C., Wahl C., Froguel P.: Routine mutation screening of HNF-1\alpha and GCK genes in MODY diagnosis: How effective are the techniques of DHPLC and direct sequencing used in combination? (Short communication) 775

Brunner G. A., Balent B., Ellmerer M., Schaupp L., Siebenhofer A., Jendle J. H., Okikawa J., Pieber T. R.: Doseresponse relation of liquid aerosol inhaled insulin in Type I diabetic patients (Short communication) 305

Bruno G., Merletti F., Biggeri A., Cerutti F., Grosso N., De Salvia A., Vitali E., Pagano G., Piemont Study Group for Diabetes Epidemiology: Increasing trend of Type I diabetes in children and young adults in the province of Turin (Italy). Analysis of age, period and birth cohort effects from 1984 to 1996 22

Byberg L., Zethelius B., McKeigue P.M., Lithell H.O.: Changes in physical activity are associated with changes in metabolic cardiovascular risk factors 2134

Cai T., Krause M. W., Odenwald W. F., Toyama R., Notkins A. L.: The IA-2 gene family: homologs in Caenorhabditis elegans, Drosophila and zebrafish 81

Cameron N. E., Cotter M. A.: Effects of an extracellular metal chelator on neurovascular function in diabetic rats 621

Cameron N. E., Tuck Z., McCabe L., Cotter M. A.: Effects of the hydroxyl radical scavenger, dimethylthiourea, on peripheral nerve tissue perfusion, conduction velocity and nociception in experimental diabetes 1161

Carlsson M., Orho-Melander M., Hedenbro J., Groop L. C.: Common variants in the β2-(Gln27Glu) and β3-(Trp64Arg)-adrenoceptor genes are associated with elevated serum NEFA concentrations and Type II diabetes 629

Carpentier A., Giacca A., Lewis G.F.: Effect of increased plasma non-esterified fatty acids (NEFAs) on argininestimulated insulin secretion in obese humans 1989

Castaing M., Péault B., Basmaciogullari A., Casal I., Czernichow P., Scharfmann R.: Blood glucose normalization upon transplantation of human embryonic pancreas into beta-cell-deficient SCID mice 2066

Cederberg J., Basu S., Eriksson U.J.: Increased rate of lipid peroxidation and protein carbonylation in experimental diabetic pregnancy 766

diabetic pregnancy 766
Ceriello A., Mercuri F., Quagliaro L.,
Assaloni R., Motz E., Tonutti L., Taboga C.: Detection of nitrotyrosine in
the diabetic plasma: evidence of oxidative stress 834

Cha B.S., Ciaraldi T.P., Carter L., Nikoulina S.E., Mudaliar S., Mukherjee R., Paterniti Jr. J.R., Henry R.R.: Peroxisome Proliferator-Activated Receptor (PPAR) γ and Retinoid X Receptor (RXR) agonists have complementary effects on glucose and lipid metabolism in human skeletal muscle 444

Chen M.-C., Proost P., Gysemans C., Mathieu C., Eizirik D. L.: Monocyte chemoattractant protein-1 is expressed in pancreatic islets from prediabetic NOD mice and in interleukin-1β-exposed human and rat islet cells 325

Choi B. C. K., Shi F.: Risk factors for diabetes mellitus by age and sex: results of the National Population Health Sur-

vev 1221

Clarke P., Gray A., Adler A., Stevens R., Raikou M., Cull C., Stratton I., Holman R on behalf of the UKPDS Group: Cost-effectiveness analysis of intensive blood-glucose control with metformin in overweight patients with Type II diabetes (UKPDS No. 51) 298

Clémenceau B., Jégou D., Martignat L., Saï P.: Long-term follow-up failed to detect in vitro transmission of fulllength porcine endogenous retroviruses from specific pathogen-free pig islets to human cells 2044

Dabrowski M., Wahl P., Holmes W. E., Ashcroft F. M.: Effect of repaglinide on cloned beta cell, cardiac and smooth muscle types of ATP-sensitive potassium channels 747

D'Amico M., Marfella R., Nappo F., Di Filippo C., De Angelis L., Berrino L., Rossi F., Giugliano D.: High glucose induces ventricular instability and increases vasomotor tone in rats 464

De Angelis L., Marfella M. A., Siniscalchi M., Marino L., Nappo F., Giugliano F., De Lucia D., Giugliano D.: Erectile and endothelial dysfunction in Type II diabetes: a possible link 1155

Dogra G., Rich L., Stanton K., Watts G. F.: Endothelium-dependent and independent vasodilation studied at normoglycaemia in Type I diabetes mellitus with and without microalbuminuria

Ek J., Andersen G., Urhammer S. A., Gæde P. H., Drivsholm T., Borch-Johnsen K., Hansen T., Pedersen O.: Mutation analysis of peroxisome proliferator-activated receptor-γ coactivator-1 (PGC-1) and relationships of identified amino acid polymorphisms to Type II diabetes mellitus 2220

Ek J., Andersen G., Urhammer S. A., Hansen L., Carstensen B., Borch-Johnsen K., Drivsholm T., Berglund L., Hansen T., Lithell H., Pedersen O.: Studies of the Pro12Ala polymorphism of the peroxisome proliferator-activated receptor-γ2 (PPAR-γ2) gene in relation to insulin sensitivity among glucose tolerant Caucasians 1170

Fiedler M., Zierath J. R., Selén G., Wallberg-Henriksson H., Liang Y., Sakariassen K. S.: 5-aminoimidazole-4-car-

boxy-amide-1-β-D-ribofuranoside treatment ameliorates hyperglycaemia and hyperinsulinaemia but not dyslipidaemia in KKAy-CETP mice 2180

Fisher R. M., Eriksson P., Hoffstedt J., Hotamisligil G. S., Thörne A., Rydén M., Hamsten A., Arner P.: Fatty acid binding protein expression in different adipose tissue depots from lean and obese individuals 1268

Forbes J. M., Soulis T., Thallas V., Panagiotopoulos S., Long D. M., Vasan S.,
 Wagle D., Jerums G., Cooper M. E.:
 Renoprotective effects of a novel inhibitor of advanced glycation 108

Garduño E., Nogues M., Merino J. M., Gutiérrez-Merino C., Henao F.: The content of glycogen phosphorylase and glycogen in preparations of sarcoplasmic reticulum-glycogenolytic complex is enhanced in diabetic rat skeletal muscle 1238

Giannattasio C., Failla M., Grappiolo A., Gamba P.L., Paleari F., Mancia G.: Progression of large artery structural and functional alterations in Type I diabetes 203

Gogos C. A., Giali S., Paliogianni F., Dimitracopoulos G., Bassaris H. P., Vagenakis A. G.: Interleukin-6 and C-reactive protein as early markers of sepsis in patients with diabetic ketoacidosis or hyperosmosis 1011

Gopaul N. K., Manraj M. D., Hébé A., Lee Kwai Yan S., Johnston A., Carrier M.J., Änggård E. E.: Oxidative stress could precede endothelial dysfunction and insulin resistance in Indian Mauritians with impaired glucose metabolism 706

Gragnoli C., Cockburn B. N., Chiaramonte F., Gorini A., Marietti G., Marozzi G.Signorini A. M.: Early-onset Type II diabetes mellitus in Italian families due to mutations in the genes encoding hepatic nuclear factor 1α and glucokinase 1326

Grunberger G., Qiang X., Li Z., Mathews S. T., Sbrissa D., Shisheva A., Sima A. A. F.: Molecular basis for the insulinomimetic effects of C-peptide 1247

Gysemans C. A., Pavlovic D., Bouillon R., Eizirik D. L., Mathieu C.: Dual role of interferon-γ signalling pathway in sensitivity of pancreatic beta cells to immune destruction 567

Hajduch E., Balendran A., Batty I.H., Litherland G.J., Blair A.S., Downes C.P., Hundal H.S.: Ceramide impairs the insulin-dependent membrane recruitment of Protein Kinase B leading to a loss in downstream signalling in L6 skeletal muscle cells 173

Hansen L., Jensen J. N., Ekstrøm C. T., Vestergaard H., Hansen T., Pedersen O.: Studies of variability in the PTEN gene among Danish caucasian patients with Type II diabetes mellitus 237 Hansen T., Ambye L., Grarup N., Hansen L., Echwald S.M., Ferrer J., Pedersen O.: Genetic variability of the *SUR1* promoter in relation to beta-cell function and Type II diabetes mellitus 1330

Hashim Y., Shepherd D., Wiltshire S., Holman R. R., Levy J. C., Clark A., Cull C. A.: Butyrylcholinesterase K variant on chromosome 3q is associated with Type II diabetes in white Caucasian subjects 2227

Heald A. H., Cruickshank J. K., Riste L. K., Cade J. E., Anderson S., Greenhalgh A., Sampayo J., Taylor W., Fraser W., White A., Gibson J. M.: Close relation of fasting insulin-like growth factor binding protein-1 (IGFBP-1) with glucose tolerance and cardiovascular risk in two populations 333

Hickner R. C., Brunson M. A., McCammon M., Mahar M. T., Garry J. P., Houmard J. A.: Diabetic groups as defined by ADA and NDDG criteria have a similar aerobic capacity, blood pressure and body composition 26

Hill C., Logan A., Smith C., Grønbaek H., Flyvbjerg A.: Angiotensin converting enzyme inhibitor suppresses glomerular transforming growth factor β receptor expression in experimental diabetes in rats 495

Hounsom L., Corder R., Patel J., Tomlinson D. R.: Oxidative stress participates in the breakdown of neuronal phenotype in experimental diabetic neuropathy 424

Huck C., Endl J., Walk T., Noessner E., Jung G., Wank R., Schendel D.J.: HLA-DR53 molecules restrict glutamic acid decarboxylase peptide presentation to T cells of a Type I diabetes patient: specification of the trimolecular HLA-peptide/T-cell receptor complex 70

Hui Y. Y., McAmis W. C., Baynes J. W., Schaeffer Jr. R. C., Wolf M. B.: Effect of advanced glycation end products on oxidative stress in endothelial cells in culture: a warning on the use of cells studied in serum-free media 1310

Isomaa B., Henricsson M., Almgren P., Tuomi T., Taskinen M.-R., Groop L.: The metabolic syndrome influences the risk of chronic complications in patients with Type II diabetes 1148

Jensen J. N., Hansen L., Ekstrøm C. T., Pociot F., Nerup J., Hansen T., Pedersen O.: Polymorphisms in the neurogenin 3 gene (*NEUROG*) and their relation to altered insulin secretion and diabetes in the Danish Caucasian population 123

Jetton T.L., Liu Y.Q., Trotman W.E., Nevin P.W., Sun X.-J., Leahy J.L.: Enhanced expression of insulin receptor substrate-2 and activation of protein kinase B/Akt in regenerating pancreatic duct epithelium of 60%-partial pancreatectomy rats 2056

Jones S. E., Bilous R. W., Flyvbjerg A., Marshall S. M.: Intra-uterine environment influences glomerular number and the acute renal adaptation to experimental diabetes 721

Jude E. B., Selby P. L., Burgess J., Lilleystone P., Mawer E. B., Page S. R., Donohoe M., Foster A. V. M., Edmonds M. E., Boulton A. J. M.: Bisphosphonates in the treatment of Charcot neuroarthropathy: a doubleblind randomised controlled trial 2032

Kaburagi Y., Satoh S., Yamamoto-Honda R., Ito T., Ueki K., Akanuma Y., Sekihara H., Kimura S., Kadowaki T.: Insulin-independent and wortmanninresistant targeting of IRS-3 to the plasma membrane via its pleckstrin homology domain mediates a different interaction with the insulin receptor from that of IRS-1 992

Karlsson E., Sandler S.: Islet amyloid polypeptide promotes beta-cell proliferation in neonatal rat pancreatic islets (Short communication) 1015

Karlsson M. G. E., Garcia J., Ludvigsson J.: Cows' milk proteins cause similar Th1- and Th2-like immune response in diabetic and healthy children 1140

Kaser S., Sandhofer A., Föger B., Ebenbichler C. F., Igelseder B., Malaimare L., Paulweber B., Patsch J. R.: Influence of obesity and insulin sensitivity on phospholipid transfer protein activity 1111

Kautzky-Willer A., Pacini G., Tura A., Bieglmayer C., Schneider B., Ludvik B., Prager R., Waldhäusl W.: Increased plasma leptin in gestational diabetes 164

Kellerer M., Lammers R., Fritsche A., Strack V., Machicao F., Borboni P., Ullrich A., Häring H. U.: Insulin inhibits leptin receptor signalling in HEK293 cells at the level of janus kinase-2: a potential mechanism for hyperinsulinaemia-associated leptin resistance 1125

Kim B.-M., Han Y.-M., Shin Y.-J., Min B.-H., Park I.-S.: Clusterin expression during regeneration of pancreatic islet cells in streptozotocin-induced diabetic rats 2192

Kimpimäki T., Erkkola M., Korhonen S., Kupila A., Virtanen S. M., Ilonen J., Simell O., Knip M.: Short-term exclusive breastfeeding predisposes young children with increased genetic risk of Type I diabetes to progressive beta-cell autoimmunity (Rapid communication) 63

Kiss G., Somogyi J., Csermely P., Szelényi J., Vér A: Streptozotocin-induced diabetes alters the oligomerization pattern of acetylcholinesterase in rat skeletal muscle (Short communication) 220 Kjems L. L., Vølund A., Madsbad S.: Quantification of beta-cell function during IVGTT in Type II and non-diabetic subjects: assessment of insulin secretion by mathematical methods 1339

Knuuti J., Takala T.O., Någren K., Sipilä H., Turpeinen A.K., Uusitupa M.I.J., Nuutila P.: Myocardial fatty acid oxidation in patients with impaired glu-

cose tolerance 184
Koopmans S. J., Jong M. C., Que I.,
Dahlmans V. E. H., Pijl H., Radder
J. K., Frölich M., Havekes L. M.: Hyperlipidaemia is associated with increased insulin-mediated glucose metabolism, reduced fatty acid metabolism and normal blood pressure in
transgenic mice overexpressing human
apolipoprotein C1 437

Kostyuk E., Voitenko N., Kruglikov I., Shmigol A., Shishkin V., Efimov A., Kostyuk P.: Diabetes-induced changes in calcium homeostasis and the effects of calcium channel blockers in rat and mice nociceptive neurons 1302

Kowluru A.: Adenine and guanine nucleotide-specific succinyl-CoA synthetases in the clonal beta-cell mitochondria: implications in the beta-cell highenergy phosphate metabolism in relation to physiological insulin secretion 89

Kristinsson S. Y., Thorolfsdottir E. T., Talseth B., Steingrimsson E., Thorsson A. V., Helgason T., Hreidarsson A. B., Arngrimsson R.: MODY in Iceland is associated with mutations in *HNF-1a* and a novel mutation in *NeuroD1* 2098

Kupila A., Muona P., Simell T., Arvilommi P., Savolainen H., Hämäläinen A.-M., Korhonen S., Kimpimäki T., Sjöroos M., Ilonen J., Knip M., Simell O.: Feasibility of genetic and immunological prediction of Type I diabetes in a population-based birth cohort 290

Kuzuya M., Asai T., Kanda S., Maeda K., Cheng X. W., Iguchi A.: Glycation cross-links inhibit matrix metalloproteinase-2 activation in vascular smooth muscle cells cultured on collagen lattice (Short communication) 433

Lally F.J., Ratcliff H., Bone A.J.: Apoptosis and disease progression in the spontaneously diabetic BB/S Rat 320

Lao T.T., Ho L.F.: α-Thalassaemia trait and gestational diabetes mellitus in Hong Kong 966

Lawrence C. L., Proks P., Rodrigo G. C., Jones P., Hayabuchi Y., Standen N. B., Ashcroft F. M.: Gliclazide produces high-affinity block of K_{ATP} channels in mouse isolated pancreatic beta cells but not rat heart or arterial smooth muscle cells 1019

Lee W.J., Lee H. W., Palmer J. P., Park K. S., Lee H. K., Park J. Y., Hong S. K., Lee K.-U.: Islet cell autoimmunity and mitochondrial DNA mutation in Korean subjects with typical and atypical Type I diabetes 2187

Lenzen S., Tiedge M., Elsner M., Lortz S., Weiss H., Jörns A., Klöppel G., Wedekind D., Prokop C.-M., Hedrich H.J.: The LEW.1AR1/Ztm-iddm rat: a new model of spontaneous insulin-dependent diabetes mellitus 1189

Leunda-Casi A., De Hertogh R., Pampfer S.: Decreased expression of fibroblast growth factor-4 and associated dysregulation of trophoblast differentiation in mouse blastocysts exposed to high D-glucose in vitro 1318

Leviev I., Kalix B., Brulhart Meynet M-Cl, James R. W.: The paraoxonase *PON1* promoter polymorphism C(-107)T is associated with increased serum glucose concentrations in non-diabetic patients 1177

Levin K., Daa Schroeder H., Alford F.P., Beck-Nielsen H.: Morphometric documentation of abnormal intramyocellular fat storage and reduced glycogen in obese patients with Type II diabetes 824

Liu Y. Q., Montanya E., Leahy J. L.: Increased islet DNA synthesis and glucose-derived lipid and amino acid production in association with betacell hyperproliferation in normoglycaemic 60% pancreatectomy rats 1026

Lohmann T., Kellner K., Verlohren H.-J., Krug J., Steindorf J., Scherbaum W. A., Seissler J.: Titre and combination of ICA and autoantibodies to glutamic acid decarboxylase discriminate two clinically distinct types of latent autoimmune diabetes in adults (LADA) 1005

Luksch A., Polak K., Matulla B., Dallinger S., Kapiotis S., Rainer G., Wolzt M., Schmetterer L.: Glucose and insulin exert additive ocular and renal vasodilator effects on healthy humans 95

Luo T. H., Zhao Y., Li G., Yuan W. T., Zhao J. J., Chen J. L., Huang W., Luo M.: A genome-wide search for Type II diabetes susceptibility genes in Chinese Hans 501

Manraj M., Francke S., Hébé A., Ramjuttun U.S., Froguel P.: Genetic and environmental nature of the insulin resistance syndrome in Indo-Mauritian subjects with premature coronary heart disease: contribution of β3-adrenoreceptor gene polymorphism and beta blockers on triglyceride and HDL concentrations 115

Marčiulionyte D., Williams A.J.K., Bingley P.J., Urbonaite B., Gale E.A.M.: A comparison of the prevalence of islet autoantibodies in children from two countries with differing incidence of diabetes 16 Massa O., Meschi F., Cuesta-Munoz A., Caumo A., Cerutti F., Toni S., Cherubini V., Guazzarotti L., Sulli N., Matschinsky F.M., Lorini R., Iafusco D., Barbetti F and the Diabetes Study Group of the Italian Society of Paediatric Endocrinology and Diabetes (SIEDP): High prevalence of glucokinase mutations in Italian children with MODY. Influence on glucose tolerance, first-phase insulin response, insulin sensitivity and BMI 898

Matsubara A., Wasson J. C., Donelan S. S., Welling C. M., Glaser B., Permutt M. A.: Isolation and characterization of the human AKTI gene, identification of 13 single nucleotide polymorphisms (SNPs), and their lack of association with Type II diabetes (Short communication) 910

Mavri A., Alessi M.C., Bastelica D., Geel-Georgelin O., Fina F., Sentocnik J. T., Stegnar M., Juhan-Vague I.: Subcutaneous abdominal, but not femoral fat expression of plasminogen activator inhibitor-1 (PAI-1) is related to plasma PAI-1 levels and insulin resistance and decreases after weight loss 2025

Meier M.-R., Desrosiers J., Bourassa P., Blaszczyk J.: Effect of Type II diabetic peripheral neuropathy on gait termination in the elderly 585

Mifsud S. A., Allen T.J., Bertram J.F., Hulthen U.L., Kelly D.J., Cooper M. E., Wilkinson-Berka J. L., Gilbert R.E.: Podocyte foot process broadening in experimental diabetic nephropathy: amelioration with renin-angiotensin blockade 878

Miyazaki Y., Glass L., Triplitt C., Matsuda M., Cusi K., Mahankali A., Mahankali S., Mandarino L.J., DeFronzo R. A.: Effect of rosiglitazone on glucose and non-esterified fatty acid metabolism in Type II diabetic patients

Mori H., Okazawa H., Iwamoto K., Maeda E., Hashiramoto M., Kasuga M.: A polymorphism in the 5' untranslated region and a Met229→ Leu variant in exon 5 of the human UCP1 gene are associated with susceptibility to Type II diabetes mellitus (Short communication) 373

Morishige N., Chikama T.-I., Sassa Y., Nishida T.: Abnormal light scattering detected by confocal biomicroscopy at the corneal epithelial basement membrane of subjects with Type II diabetes

Mutus B., Rabini R. A., Staffolani R., Ricciotti R., Fumelli P., Moretti N., Martarelli D., Mazzanti L.: Homocysteine-induced inhibition of nitric oxide production in platelets: a study on healthy and diabetic subjects (Short communication) 979

Nagisa Y., Shintani A., Nakagawa S.: The angiotensin II receptor antagonist candesartan cilexetil (TCV-116) ameliorates retinal disorders in rats 883

Nakamura J., Kasuva Y., Hamada Y., Nakashima E., Naruse K., Yasuda Y., Kato K., Hotta N.: Glucose-induced hyperproliferation of cultured rat aortic smooth muscle cells through polyol pathway hyperactivity 480

Nakhoul F.M., Zoabi R., Kanter Y., Zoabi M., Skorecki K., Hochberg I., Leibu R., Miller B., Levy A. P.: Haptoglobin phenotype and diabetic nephropathy (Short communication) 602

Nicoletti F., Conget I., Di Marco R., Speciale A. M., Morinigo R., Bendtzen K., Gomis R.: Serum levels of the interferon-γ-inducing cytokine interleukin-18 are increased in individuals at high risk of developing Type I diabetes (Rapid communication) 309

Nordt T.K., Bode C., Sobel B.E.: Stimulation in vivo of expression of intraabdominal adipose tissue plasminogen activator inhibitor Type I by proinsu-

Not T., Tommasini A., Tonini G., Buratti E., Pocecco M., Tortul C., Valussi M., Crichiutti G., Berti I., Trevisiol C., Azzoni E., Neri E., Torre G., Martelossi S., Soban M., Lenhardt A., Cattin L., Ventura A.: Undiagnosed coeliac disease and risk of autoimmune disorders in subjects with Type I diabetes mellitus 151

Obrosova I.G., Minchenko A.G., Marinescu V., Fathallah L., Kennedy A., Stockert C. M., Frank R. N., Stevens M.J.: Antioxidants attenuate early up regulation of retinal vascular endothelial growth factor in streptozotocindiabetic rats 1102

Okada T., Tobe K., Hara K., Yasuda K., Kawaguchi Y., Ikegami H., Ito C., Kadowaki T.: Variants of neurogenin 3 gene are not associated with Type II diabetes in Japanese subjects 241

Osicka T. M., Kiriazis Z., Pratt L. M., Jerums G., Comper W.D.: Ramipril and aminoguanidine restore renal lysosomal processing in streptozotocin diabetic rats 230

Osmond D. T. D., King R. G., Brennecke S. P., Gude N. M.: Placental glucose transport and utilisation is altered at term in insulin-treated, gestationaldiabetic patients 1133

Palmén T., Ahola H., Palgi J., Aaltonen P., Luimula P., Wang S., Jaakkola I., Knip M., Otonkoski T., Holthöfer H.: Nephrin is expressed in the pancreatic beta cells 1274

Panerai A. E., Nicoletti F., Sacedote P., Arvidsson L., Conget I., Gomis R., Bartorelli A., Sandler S.: MFP14, a multifunctional emerging protein with immunomodulatory properties, prevents spontaneous and recurrent autoimmune diabetes in NOD mice 839

Pérez-Jiménez F., López-Miranda J., Pinillos M. D., Gómez P., Paz-Rojas E, Montilla P., Marín C., Velasco M. J., Blanco-Molina A., Jiménez Perepérez J. A., Ordovás J. M.: A Mediterranean and a high-carbohydrate diet improve glucose metabolism in healthy young persons 2038

Perrott R. L., North R. V., Drasdo N., Ahmed K. A., Owens D. R.: The influence of plasma glucose upon pulsatile ocular blood flow in subjects with Type II diabetes mellitus 700

Pessler D., Rudich A., Bashan N.: Oxidative stress impairs nuclear proteins binding to the insulin responsive element in the GLUT4 promoter 2156

Petersen K. F., Sullivan J. T.: Effects of a novel glucagon receptor antagonist (Bay 27-9955) on glucagon-stimulated glucose production in humans 2018

Pierucci D., Cicconi S., Bonini P., Ferrelli F., Pastore D., Matteucci C., Marselli L., Marchetti P., Ris F., Halban P., Oberholzer J., Federici M., Cozzolino F., Lauro R., Borboni P., Marlier L. N. J. L.: NGF-withdrawal induces apoptosis in pancreatic beta cells in vitro 1281

Pinizzotto M., Castillo E., Fiaux M., Temler E., Gaillard R. C., Ruiz J.: Paraoxonase2 polymorphisms are associated with nephropathy in Type II diabetes (Short communication) 104

Porta M., Sjoelie A.-K., Chaturvedi N., Stevens L., Rottiers R., Veglio M., Fuller J. H., and the EURODIAB Prospective Complications Study Group: Risk factors for progression to proliferative diabetic retinopathy in the EURODIAB Prospective Complications Study 2203

Poulsen P., Vaag A., Kyvik K., Beck-Nielsen H.: Genetic versus environmental aetiology of the metabolic syndrome among male and female twins 537

Price R. A., Li W.-D., Bernstein A., Crystal A., Golding E.M., Weisberg S.J., Zuckerman W.A.: A locus affecting obesity in human chromosome region 10p12 (Short communication) 363

Radziuk J., Pye S.: Production and metabolic clearance of glucose under basal conditions in Type II (non-insulin-dependent) diabetes mellitus (Rapid communication) 983

Ragnarson Tennvall G., Apelqvist J.: Prevention of diabetes-related foot ulcers and amputations: a cost-utility analysis based on Markov model simulations 2077

Ramachandran A., Snehalatha C., Kapur A., Vijay V., Mohan V., Das A. K., Rao P. V., Yajnik C. S., Prasanna Kumar

K. M., Nair JD for the Diabetes Epidemiology Study Group in India (DESI): High prevalence of diabetes and impaired glucose tolerance in India: National Urban Diabetes Survey 1094

Redondo M.J., Yu L., Hawa M., Mackenzie T., Pyke D. A., Eisenbarth G.S., Leslie R.D.G.: Heterogeneity of Type I diabetes: analysis of monozygotic twins in Great Britain and the United States 354

Rieusset J., Chambrier C., Bouzakri K., Dusserre E., Auwerx J., Riou J.-P., Laville M., Vidal H.: The expression of the p85α subunit of phosphatidylinositol 3-Kinase is induced by activation of the peroxisome proliferator-activated receptor γ in human adipocytes 544

Righetti A. E., Boer-Lima P. A., Lopes de Faria J. B.: The presence of genetic hypertension stimulates early renal accumulation of fibronectin in experimental diabetes mellitus (Short communi-

cation) 2088

Rigla M., Fontcuberta J., Mateo J., Caixàs A., Pou J. M., de Leiva A., Pérez A.: Physical training decreases plasma thrombomodulin in Type I and Type II diabetic patients 693

Rosenstock M., Greenberg A.S., Rudich A.: Distinct long-term regulation of glycerol and non-esterified fatty acid release by insulin and TNF-α in 3T3-

L1 adipocytes 55

Rosmond R., Chagnon M., Bouchard C., Björntorp P.: A missense mutation in the human melanocortin-4 receptor gene in relation to abdominal obesity

and salivary cortisol 1335

Sadeharju K., Lönnrot M., Kimpimäki T., Savola K., Erkkilä S., Kalliokoski T., Savolainen P., Koskela P., Ilonen J., Simell O., Knip M., Hyöty H.: Enterovirus antibody levels during the first two years of life in prediabetic autoantibody-positive children 818

Sasaoka T., Hori H., Wada T., Ishiki M., Haruta T., Ishihara H., Kobayashi M.: SH2-containing inositol phosphatase 2 negatively regulates insulin-induced glycogen synthesis in L6 myotubes

1258

Saukkonen T., Ilonen J., Åkerblom H. K., Savilahti E and the Childhood Diabetes in Finland Study Group: Prevalence of coeliac disease in siblings of patients with Type I diabetes is related to the prevalence of DQB1*02 allele 1051

Schoenle E.J., Lang-Muritano M., Gschwend S., Laimbacher J., Mullis P.E., Torresani T., Biason-Lauber A., Molinari L.: Epidemiology of Type I diabetes mellitus in Switzerland: steep rise in incidence in under 5 year old children in the past decade 286

Schultz C.J., Dalton R.N., Neil H.A.W., Konopelska-Bahu T., Dunger DB on behalf of the Oxford Regional Prospective Study Group: Markers of renal tubular dysfunction measured annually do not predict risk of microalbuminuria in the first few years after diagnosis of Type I diabetes 224

Seino S on behalf of The Study Group of Comprehensive Analysis of Genetic Factors in Diabetes Mellitus: S2OG mutation of the amylin gene is associated with Type II diabetes in Japanese (Short communication) 906

(Short communication) 906 Service F.J., O'Brien P.C.: The relation of glycaemia to the risk of development and progression of retinopathy in the Diabetic Control and Complica-

tions Trial 1215

Sima A. A. F., Zhang W., Sugimoto K., Henry D., Li Z., Wahren J., Grunberger G.: C-peptide prevents and improves chronic Type I diabetic polyneuropathy in the BB/Wor rat 889

Stallmeyer B., Pfeilschifter J., Frank S.: Systemically and topically supplemented leptin fails to reconstitute a normal angiogenic response during skin repair

in diabetic ob/ob mice 471

Stingl H., Krššak M., Krebs M., Bischof M.G., Nowotny P., Fürnsinn C., Shulman G.I., Waldhäusl W., Roden M.: Lipid-dependent control of hepatic glycogen stores in healthy humans 48

Stratton I. M., Kohner E. M., Aldington S. J., Turner R. C., Holman R. R., Manley S. E., Matthews DR for the UKPDS Group: UKPDS 50: Risk factors for incidence and progression of retinopathy in Type II diabetes over 6 years from diagnosis 156

Takeda M., Mori F., Yoshida A., Takamiya A., Nakagomi S., Sato E., Kiyama H.: Constitutive nitric oxide synthase is associated with retinal vascular permeability in early diabetic rats 1043

Tatarkiewicz K., Garcia M., Omer A., Van Schilfgaarde R., Weir G.C., De Vos P.: C-Peptide responses after meal challenge in mice transplanted with microencapsulated rat islets 646

Thams P., Capito K.: Differential mechanisms of glucose and palmitate in augmentation of insulin secretion in mouse pancreatic islets 738

Torbjörnsdotter T.B., Jaremko G.A., Berg U.B.: Ambulatory blood pressure and heart rate in relation to kidney structure and metabolic control in adolescents with Type I diabetes 865

Trautner C., Haastert B., Giani G., Berger M.: Incidence of blindness in southern Germany between 1990 and 1998 147

Tsuchida A., Nakagawa T., Itakura Y., Ichihara J., Ogawa W., Kasuga M., Taiji M., Noguchi H.: The effects of brain-derived neurotrophic factor on insulin signal transduction in the liver of diabetic mice 555

Tsunoda K., Sanke T., Nakagawa T., Furuta H., Nanjo K.: Single nucleotide polymorphism (D68D, T to C) in the syntaxin 1A gene correlates to age at onset and insulin requirement in Type II diabetic patients 2092

Ubels F. L., Muntinga J. H. J., Links T. P., Hoogenberg K., Dullaart R. P. F., Reitsma W. D., Smit A. J.: Redistribution of blood volume in Type I diabetes (Short communication) 429

Ukkola O., Sun G., Bouchard C.: Insulinlike growth factor 2 (*IGF2*) and *IGF*binding protein 1 (*IGFBP1*) gene variants are associated with overfeeding-

induced metabolic changes 2231
Vella A., Shah P., Basu R., Basu A., Camilleri M., Schwenk W.F., Rizza R. A.:
Type I diabetes mellitus does not alter initial splanchnic glucose extraction or hepatic UDP-glucose flux during enteral glucose administration 729

Vessby B., Uusitupa M., Hermansen K., Riccardi G., Rivellese A. A., Tapsell L. C., Nälsén C., Berglund L., Louheranta A., Rasmussen B. M., Calvert G. D., Maffetone A., Pedersen E., Gustafsson I.-B., Storlien L. H.: Substituting dietary saturated for monoensaturated fat impairs insulin sensitivity in healthy men and women: The KANWU study 312

Virtanen K. A., Peltoniemi P., Marjamäki P., Asola M., Strindberg L., Parkkola R., Huupponen R., Knuuti J., Lönnroth P., Nuutila P.: Human adipose tissue glucose uptake determined using [¹⁸F]-fluoro-deoxy-glucose ([¹⁸F]FDG) and PET in combination with micro-

dialysis 2171

Waterworth D. M., Talmud P.J., Humphries S. E., Wicks P.D., Sagnella G. A., Strazzullo P., Alberti K. G. M. M., Cook D. G., Cappuccio F. P.: Variable effects of the *APOC3 -482C > T* variant on insulin, glucose and triglyceride concentrations in different ethnic groups 245

Weets I., Van Autreve J., Van der Auwera B.J., Schuit F.C., Du Caju M. V. L., Decochez K., De Leeuw I. H., Keymeulen B., Mathieu C., Rottiers R., Dorchy H., Quartier E., Gorus FK and the Belgian Diabetes Registry: Male-to-female excess in diabetes diagnosed in early adulthood is not specific for the immune-mediated form nor is it HLA-DQ restricted: possible relation to increased body mass index 40

Weiss U., Cervar M., Puerstner P., Schmut O., Haas J., Mauschitz R., Arikan G., Desoye G.: Hyperglycaemia in vitro alters the proliferation and mitochondrial activity of the choriocarcinoma cell lines BeWo, JAR and JEG-3 as models for human first-trimester trophoblast 209 Weng J., Macfarlane W. M., Lehto M., Gu H. F., Shepherd L. M., Ivarsson S. A., Wibell L., Smith T., Groop L. C.: Functional consequences of mutations in the MODY4 gene (IPF1) and coexistence with MODY 3 mutations 249

Wice B. M., Bernal-Mizrachi E., Permutt M. A.: Glucose and other insulin secretagogues induce, rather than inhibit, expression of *Id-1* and *Id-3* in pancreatic islet beta cells 453

Wolford J. K., Hanson R. L., Bogardus C., Prochazka M.: Analysis of the *Lamin A/C* gene as a candidate for Type II diabetes susceptibility in Pima Indians 779

Wu G., Haynes T. E., Yan W., Meininger C.J.: Presence of glutamine: fructose-6-phosphate amidotransferase for glucosamine-6-phosphate synthesis in endothelial cells: effects of hyperglycaemia and glutamine 196

Wu L., Eftekharpour E., Davies G.F., Roesler W.J., Juurlink B.H.J.: Troglitazone selectively inhibits glyoxalase I

gene expression 2004

Yu H. Y., Inoguchi T., Kakimoto M., Nakashima N., Imamura M., Hashimoto T., Umeda F., Nawata H.: Saturated non-esterified fatty acids stimulate de novo diacylglycerol synthesis and protein kinase C activity in cultured aortic smooth muscle cells 614

Zhang D., Kaufman P.L., Gao G., Saunders R. A., Ma J. X.: Intravitreal injection of plasminogen kringle 5, an endogenous angiogenic inhibitor, arrests retinal neovascularization in rats 757

Zhong Z.-H., Pramanik A., Ekberg K., Jansson O. T., Jörnvall H., Wahren J., Rigler R.: Insulin binding monitored by fluorescence correlation spectroscopy 1184

Zilberfarb V., Siquier K., Strosberg A. D., Issad T.: Effect of dexamethasone on adipocyte differentiation markers and tumour necrosis factor-a expression in human PAZ6 cells 377

For debate

Airaksinen K. E. J.: Silent coronary artery disease in diabetes – a feature of autonomic neuropathy or accelerated atherosclerosis? 259

Kolb H., Gale E. A. M.: Does partial preservation of residual beta-cell function justify immune intervention in recent onset Type I diabetes? 1349

Wilkin T.J.: The accelerator hypothesis: weight gain as the missing link between Type I and Type II diabetes 914

Letters

Albareda M.: et al. (2000) Assessment of insulin sensitivity and beta-cell function from measurements in the fasting state and during an oral glucose tolerance test. Diabetologia 43: 1507–1511 783

Allcock R., Cheong K., Christiansen F., Witt C.: Comments to: Gambelunghe G., Ghaderi, Cosentino A et al. (2000) Association of MHC Class I hainrelated A (MIC-A) gene polymorphsm with Type I diabetes. Diabetologia 43: 507-514 514

Benhamou P. Y.: et al. (2001) Human islet transplantation network for the treatment of Type I diabetes: first data from the Swiss-French GRAGIL consortium (1999–2000). Diabetologia 44: 859–864 2239

Bergman RN (2000) Non-esterified fatty acids and the liver: why is insulin secreted into the portal vein? Diabetologia 43: 946–952 784

Boomsma F., Ekberg K., Bruining G.J.: C Peptide and insulin do not influence plasma semicarbazide-sensitive amine oxidase activity 388

Cavallo M. G., Romeo S., Coppolino G., Pozzilli P.: Continuous glucose monitoring during the European Soccer cup semifinal, Italy against Holland 268

Charlton R., Smith G., Day A.: Munchausen's syndrome manifesting as factitious hypoglycaemia 784

Dalgaard L. T., Hansen T., Urhammer S. A., Drivsholm T., Borch-Johnsen K., Pedersen O.: The uncoupling protein 3-55 C -> T variant is not associated with Type II diabetes mellitus in Danish subjects 1065

Ebeling P., Heliövaara M. K.: Comments to: Yudkin J, Panahloo A, Stenhouwer C et al. (2000) The influence of improved glycaemic control with insulin and sulphonylureas on acute phase and endothelial markers in Type II diabetic subjects. Diabetologia 43: 1099–1106 519

Fernandez-Real J.-M., Ricart W.: Comments to: Yudkin J, Panahloo A, Stenhouwer C et al. (2000) The influence of improved glycaemic control with insulin and sulphonylureas on acute phase and endothelial markers in Type II diabetic subjects. Diabetologia 43: 1099-1106 518

Gambelunghe G., Falorni A., Sanjeevi C. B.: The authors' reply 515

Gloyn A. L., Ashcroft S. J. H.: The betacell Ca²⁺/calmodulin-dependent protein kinase II (CaM kinase II) β3 isoform containing a proline-rich tandem repeat in the association domain can be found in the human genome 787

Harder T., Plagemann A.: Comments to: Velho G, Hattersley AT, Froguel P (2000) Maternal diabetes alters birth weight in glucokinase- deficient (MODY2) kindred but has no influence on adult weight, height, insulin secretion or insulin sensitivity Diabetologia 43: 1060–1063 516

Icks A., Rosenbauer J., Holl R.W., Grabert M., Giani G on behalf of the German Working Group on Quality Control in Childhood Diabetes: Increased hospitalization with longer distance from treatment centre in diabetic pediatric patients in Germany 1068

Iwasaki N., Babazono T., Tomonaga O., Ogata M., Yokokawa H., Iwamoto Y.: Mutations in the hepatocyte nuclear factor-1ß (MODY5) gene are not a major factor contributing to end-stage renal disease in Japanese people with diabetes mellitus 127

Iwasaki N., Okabe I., Momoi M. Y., Ohashi H., Ogata M., Iwamoto Y.: Splice site mutation in the hepatocyte nuclear factor-1β Gene, IVS2nt + 1G > A, associated with maturity-onset diabetes of the young, renal dysplasia and bicornuate uterus 387

Iwata E., Matsuda H., Fukuda T., Fukuen S., Motomura T., Igarashi T., Yamamoto I., Azuma J.: Mutations of the peroxisome proliferator-activated receptor γ ($PPAR\gamma$) gene in a Japanese population: the Pro12Ala mutation in $PPAR\gamma$ 2 is associated with lower concentrations of serum total and non-HDL cholesterol 1354

Jones C. J. P., Weiss U., Simán C. M., Desoye G.: Mitochondria from human trophoblast and embryonic liver cells are resistant to hyperglycaemia-associated high-amplitude swelling 389

Lampasona V., Scirpoli M., Bosi E., Bonifacio E.: ICA12(SOX13) autoantibodies are unlikely to be a useful marker for pre-clinical Type I diabetes 267

Lindi V., Sivenius K., Niskanen L., Laakso M., Uusitupa M. I. J.: Effect of the *Pro12Ala* polymorphism of the *PPAR*-γ2 gene on long-term weight change in Finnish non-diabetic subjects 925

Lipinski B., Lipinska I., Kato Y.: Is the macromolecular protein complex (MPC) a marker for oxidative stress in diabetes mellitus? 1356

Maier S., Lange V., Simm A., Walter U., Kirstein M.: Insulin fails to modulate the cardiac L-Type Ca2+ current in Type II diabetes patients – a possible link to cardiac dysfunction in diabetes mellitus 269

Myers M. A., Mackay I. R., Rowley M. J., Zimmet P. Z.: Dietary microbial toxins and Type I diabetes – a new meaning for seed and soil 1199

Nakhoul F. M.: et al. (2001) Haptoglobin phenotype and diabetes. Diabetologia 44: 602–604 2237 Novials A., Rojas I., Franco C., Casamitjana R., Usac E. F., Gomis R.: A novel mutation in islet amyloid polypeptide (IAPP) gene promotor is associated with Type II diabetes mellitus 1064

Ohyama Y., Utsugi T., Uchiyama T., Hanaoka T., Hasegawa A., Tomono S., Nagai R., Kurabayashi M.: The elevated proinsulin-to-insulin ratio is associated with oxidative stress in Down's syndrome 788

Osmond D. T. D., Nolan C. J., King R. G., Brennecke S. P., Gude N. M.: Effects of gestational diabetes on human placental glucose uptake, transfer, and utilisation 927

Pinizzotto M., Castillo E., Fiaux M., Temler E., Gaillard R. C., Ruiz J.: (2001) Paraoxonase 2 polymorphisms are associated with diabetic nephropathy in Type II diabetes. Diabetologia 44: 104–107 1062

Redondo M.J., Yu L., Hawa M., Mackenzie T., Pyke D. A., Eisenbarth G.S., Leslie R. D. G.: Heterogeneity of Type I diabetes: analysis of monozygotic twins in Great Britain and the United States 928

Renard C. B., Bornfeldt K. E.: Human arterial smooth muscle cells rapidly deplete cell culture media of glucose 1067

Shield J., Owen K., Robinson D.O., Mackay D., Ellard S., Hattersley A., Temple I. K.: Maturity onset diabetes of the young (MODY) and early onset Type II diabetes are not caused by loss of imprinting at the transient neonatal diabetes (TNDM) locus 924

Sir-Petermann T., Pérez-Bravo F., Angel B., Maliqueo M., Calvillán M., Palomino A.: *G972R* polymorphism of IRS-1 in women with polycystic ovary syndrome 1200

Skoog T., Eriksson P., Hoffstedt J., Rydén M., Hamsten A., Arner P.: Tumour necrosis factor-α (TNF-α) polymorphisms –857C/A and –863C/A are associated with TNF-α secretion from human adipose tissue 654

Soltész G., Rózsai B.: Comments to: Tarnow L, Kjeld T, Knudsen E et al. (2000) Lack of synergism between long-term poor glycaemic control and three gene polymorphisms of the renin angiotensin system on risk of developing diabetic nephropathy in Type I diabetic patients. Diabetologia 43: 794–799 520

Vaxillaire M., Benmezroua Y., Durand E., Vasseur F., Froguel P.: No evidence for diabetes-associated mutations of PEK/EIF2AK3 gene in French patients with early-onset Type II diabetes 786

Velho G., Hattersley A. T., Froguel P.: The authors' reply 517

Waldhäusl W (2001) Editorial: Finally we have arrived in a new millennium.

Diabetologia 44: 1–2 923

Yajnik C.S., Coyaji K.J., Joglekar C.V., Keilingray S., Fall C.: Paternal insulin resistance and fetal growth: problem for the 'fetal insulin' and the 'fetal origins' hypotheses 1197

Book reviews

270, 1202

EASD News section

1, 13, 17, 19, 25, 27, 29, 35, 41, 55, 67

Difficult Diabetes

1202

Erratum

391

List of referees

2240

Author index

2245

Subject index

2250

Contents of Volume 44

I



